

Prescription Label

Patient Name:

Species:

Drug Name & Strength:

Directions (amount to give how often & for how long):

Prescribing Veterinarian's Name & Contact Information:

Refills:

[Content to be provided by prescribing veterinarian]

Phenylpropanolamine

(fen-ill-proe-pa-**nole**-a-meen)

Description:

Sympathomimetic

Other Names for this Medication:

PPA, Proin[®], Proin ER[®]

Common Dosage Forms:

Veterinary: 25 mg, 50 mg, & 75 mg Chewable Tablets (liver flavored, scored); 18 mg, 38 mg, 74 mg, & 145 mg Extended-Release Tablets.

Human: None.

This information sheet does not contain all available information for this medication and has not been reviewed by FDA Center for Veterinary Medicine. This sheet can help answer commonly asked questions but is not a substitute for medical advice. If you have other questions or need more information about this medication, contact your veterinarian or pharmacist.

Key Information

- Commonly prescribed to reduce dribbling of urine (urinary incontinence).
- May be given with or without food. Giving with food may help if your animal vomits or acts sick after getting the drug.
- Tablets are liver-flavored; be sure to keep out of reach of animals.

How is this medication useful?

Phenylpropanolamine is used to treat urinary incontinence in dogs and sometimes cats.

The FDA (U.S. Food & Drug Administration) has approved phenylpropanolamine chewable tablets and extended-release tablets for use in dogs. The FDA allows veterinarians to prescribe and use products containing this drug in different species or for other conditions in certain situations. You and your veterinarian can discuss why this drug is the most appropriate choice.

What should I tell my veterinarian to see if this medication can be safely given?

Many things might affect how well this drug will work in your animal. Be sure to discuss the following with your veterinarian so together you can make the best treatment decisions.

- Other drugs can interact with phenylpropanolamine, so be sure to tell your veterinarian and pharmacist what medications (including vitamins, supplements, or herbal therapies) you give your animal, including the amount and time you give each.
- Tell your veterinarian about any conditions or diseases your animal may have now or has had in the past.
- If your animal has been treated for the same disease or condition in the past, tell your veterinarian about the treatment and how well it did or didn't work.
- If your animal is pregnant or nursing, talk to your veterinarian about the risks of using this drug.
- Tell your veterinarian and pharmacist about any medication side effects (including allergic reactions, lack of appetite, diarrhea, itching, hair loss) your animal has developed in the past.

How long until I will know if this medication is working, and how long will the effects of this medication last?

This medication should help your animal feel better within 1 to 2 hours. Your animal's clinical signs should improve after that time. The effects of this medication are short-lived, meaning they will stop working within 24 hours, although the benefits may be prolonged if your animal has decreased kidney and/or liver function.

When should this medication not be used or be used very carefully?

No drug is 100% safe in all patients, but your veterinarian will discuss with you any specific concerns about using this drug in your animal.

This drug **SHOULD NOT** be used in patients:

- That are allergic to it.

This drug should be used **WITH CAUTION** in patients with:

- Glaucoma
- Seizure disorders (convulsions)
- High blood pressure
- Heart, thyroid, kidney, or liver disease

If your animal has any of these conditions, talk to your veterinarian about the potential risks versus benefits.

What are the side effects of this medication?

Common but not serious side effects include:

- Vomiting, lack of appetite, higher level of activity, restlessness, behavior changes, diarrhea, and greater thirst.

You don't have to be overly concerned if you see any of these signs unless they worsen or persist. Contact your veterinarian if this happens.

Side effects that may be serious or indicate a serious problem:

- Seizures, collapse (passing out), or stroke-like signs (eg, not able to walk)

If you see any of these signs, contact your veterinarian immediately.

If my animal gets too much of this medication (an overdose), what should I do?

Overdoses of phenylpropanolamine can be serious. If you witness or suspect an overdose, immediately contact your veterinarian or an animal poison control center for further advice. Animal poison control centers that are open 24 hours a day include: **Pet Poison HELPLINE** (855-764-7661) and **ASPCA Animal Poison Control Center** (888-426-4435); a consultation fee is charged for these services.

How should this medication be given?

For this medication to work, give it exactly as your veterinarian has prescribed. It's a good idea to always check the prescription label to be sure you are giving the drug correctly.

- Phenylpropanolamine may be given with or without food, but if your animal vomits or acts sick after receiving the drug on an empty stomach, try giving the next dose with food or a small treat. If vomiting continues, contact your veterinarian.
- Do not split or crush the extended-release tablets.
- If you have difficulty getting your animal to take the medicine, contact your veterinarian or pharmacist for tips to help dosing and reducing the stress of medication time.
- This medication can be given for various lengths of time. Be sure you understand how long your veterinarian wants you to continue giving this medication. Prescription refills may be necessary before the therapy will be complete. Before stopping this medication, talk to your veterinarian, as there may be important reasons to continue its use.

What should I do if I miss giving a dose of this medication?

If you miss a dose, give it when you remember, but if it is close to the time for the next dose, skip the dose you missed and give it at the next scheduled time. After that, return to the regular dosing schedule. Do not double-up or give extra doses.

How should I store this medication?

Store this medication in the original prescription bottle or an approved dosage reminder container (ie, pill minder) at room temperature, out of direct sunlight. Keep away from children and other animals.

Can handling this medication be hazardous to me, my family, or other animals?

There are no specific precautions required when handling this medication unless you are allergic to it. Wash your hands after handling any medication.

How should I dispose of this medication if I don't use it all?

- Do not flush this medication down the toilet or wash it down the sink. If a community drug "take-back" program is available, use this option. If there is no take-back program, mix the drug with coffee grounds or cat litter (to make it undesirable to children and animals and unrecognizable to people who might go through your trash), place the mixture in a sealable plastic bag to keep it from leaking out, and throw the bag out with the regular trash.
- Do not save leftover medication for future use or give it to others to use.

What other information is important for this medication?

- Although the chewable tablets (Proin®) are liver-flavored, they do not contain beef or beef by-products.
- Use of this drug may not be allowed in certain animal competitions. Check rules and regulations before entering your animal in a competition while this medication is being administered.

If you have any other questions about this medication, contact your veterinarian or pharmacist.

Special Instructions
